

STRONGLY EXPOSED POINTS IN BASES FOR THE POSITIVE CONE OF ORDERED BANACH SPACES AND CHARACTERIZATIONS OF $l_1(\Gamma)$

by IOANNIS A. POLYRAKIS

(Received 27th May 1985)

1. Introduction

The study of extreme, strongly exposed points of closed, convex and bounded sets in Banach spaces has been developed especially by the interconnection of the Radon–Nikodým property with the geometry of closed, convex and bounded subsets of Banach spaces [5], [2]. In the theory of ordered Banach spaces as well as in the Choquet theory, [4], we are interested in the study of a special type of convex sets, not necessarily bounded, namely the bases for the positive cone. In [7] the geometry (extreme points, dentability) of closed and convex subsets K of a Banach space X with the Radon–Nikodým property is studied and special emphasis has been given to the case where K is a base for a cone P of X . In [6, Theorem 1], it is proved that an infinite-dimensional, separable, locally solid lattice Banach space is order-isomorphic to l_1 if, and only if, X has the Krein–Milman property and its positive cone has a bounded base.

In this paper (Section 3) we study the existence of strongly exposed points in a base B for a cone P of a Banach space X and we characterize the strongly exposing functionals. It is proved (Theorem 3.1) that the existence of strongly exposed points in a base B for P is closely connected with the existence of a bounded base for the cone P .

In Section 4 we prove a similar result to [6, Theorem 1] for the space $l_1(\Gamma)$, (Theorem 4.1). Afterwards we prove that if a Banach space X ordered by the closed, generating cone P has the R.D.P. then X is order-isomorphic to $l_1(\Gamma)$ if, and only if, P has the Krein–Milman property and $\text{sep}(B) \neq 0$, for at least one base B for P , (Proposition 4.2).

2. Notations and definitions

Let X be a normed space, K a convex subset of X and $x_0 \in K$. We say that x_0 is an exposed point of K if there exists a continuous linear functional g of X ($g \in X^*$) such that $g(x_0) > g(x), \forall x \in K \setminus \{x_0\}$. In this case we say that g exposes x_0 in K . We say that x_0 is a strongly exposed point of K if there exists $g \in X^*$ which exposes x_0 in K and for each sequence (x_n) of K , $g(x_n) \rightarrow g(x_0)$ implies $x_n \rightarrow x_0$. In this case we say that g strongly exposes x_0 in K . We denote by $\text{ep}(K)$, $\text{sep}(K)$ the set of extreme, strongly exposed points of K . For each $A \subseteq X$ we denote by \bar{A} the closure of A and by $\text{c}\bar{\text{co}} A$ the closed convex hull of A . Let X be ordered by the cone P . $x \in P \setminus \{0\}$ is an external point of P .

$x \in \text{EP}(P)$, if for each $y \in P$, $0 < y < x$ implies $y = \lambda x$. A subset B of P is a base for P if there exists a strictly positive linear functional f of X such that $B = \{x \in P \mid f(x) = 1\}$. Then we say that the base B is defined by the functional f . The cone P is generating if $X = P - P$. We say that P is well-based if there exists a bounded base B for P and $0 \notin \bar{B}$. The space X is a locally solid linear lattice if X is a linear lattice and there exists a real number $a > 0$ such that for each $x, y \in X$, $|x| \leq |y|$, implies $\|x\| \leq a\|y\|$. A linear functional f of X is uniformly monotonic if there exists a real number $a > 0$ such that $f(x) \geq a\|x\|, \forall x \in P$.

It is easy to show that g strongly exposes 0 in P if, and only if, $-g$ is uniformly monotonic and therefore that $0 \in \text{sep}(P)$ if, and only if, P is well-based. X is order-isomorphic to an ordered normed space Y if there exists an isomorphism T of X onto Y and T, T^{-1} are positive. An ordered linear space X has the Riesz decomposition property (R.D.P.) if for any three positive elements x, y, z of X with $x \leq y + z$ there exist $x_1, x_2 \in X$ such that $0 \leq x_1 \leq y$, $0 \leq x_2 \leq z$ and $x = x_1 + x_2$. Each linear lattice has the R.D.P.

3. Strongly exposed points in a base for a cone

Proposition 3.1. *Let X be a normed space and K be an unbounded, convex subset of X . If g strongly exposes the point x_0 in K , then $g(x_v) \rightarrow -\infty$, for each sequence (x_v) of K with $\|x_v\| \rightarrow +\infty$.*

Proof. Let $\rho \in \mathbb{R}$ such that $\|x_0\| < \rho$ and (x_v) be a sequence of K with $\lim_{v \rightarrow \infty} \|x_v\| = +\infty$. Then there exists a sequence (y_v) of K , such that

$$\|y_v\| = \rho \text{ and } y_v = \lambda_v x_0 + (1 - \lambda_v)x_v, \quad \lambda_v \in (0, 1).$$

Then

$$\rho = \|y_v\| \geq |\lambda_v\|x_0\| - (1 - \lambda_v)\|x_v\|$$

and therefore $\lambda_v \rightarrow 1$.

If $(g(x_{k_v}))$ is a bounded subsequence of $(g(x_v))$, then

$$g(y_{k_v}) \rightarrow g(x_0)$$

and therefore $y_{k_v} \rightarrow x_0$. This is a contradiction. Hence $g(x_v) \rightarrow -\infty$ because $g(x_v) \leq g(x_0)$.

Let X be a normed space, K a convex subset of X , $u \in X^*$ such that $u(K) = \{1\}$ and $\lambda \in \mathbb{R}$. By a simple computation we have:

(S₁) A functional $g \in X^*$ strongly exposes x_0 in K if, and only if, $g - \lambda u$ strongly exposes x_0 in K .

Since $(g - g(x_0)u)(x_0) = 0$, we have that

(S₂) x_0 is a strongly exposed point of K if, and only if, there exists $g \in X^*$ such that g strongly exposes x_0 in K and $g(x_0) = 0$.

If g is as in (S_2) , then $g(x) \leq 0$ for each $x \in K$ and we shall say that g is a *negative-strongly exposing functional* of x_0 in K . Let X be ordered by the cone P and x_0 be an extremal point of P . We shall say that x_0 has a *continuous projection* P_{x_0} (or that x_0 is an extremal point of P with continuous projection P_{x_0}) if, and only if, P_{x_0} is a linear, continuous, positive projection of X onto $[x_0]$ such that $P_{x_0}(x) \leq x$, for each $x \in P$.

Let x_0 be an extremal point of P with continuous projection P_{x_0} . We denote by $\rho(x_0, \cdot)$ the continuous linear functional of X , defined by the formula

$$P_{x_0}(x) = \rho(x_0, x)x_0, \quad x \in X.$$

Also for each $h \in X^*$ we denote by h_{x_0} the functional

$$h_{x_0}(x) = h(P_{x_0}(x) - x), \quad x \in X.$$

If B is a base for P defined by $f \in X^*$ and $x'_0 = \lambda x_0 \in B$, then we have:

- (P₁) $P_{x_0}(x) < x'_0$, for each $x \in B \setminus \{x'_0\}$,
- (P₂) for each sequence (x_n) of B , $f(P_{x_0}(x_n)) \rightarrow 1$ implies $P_{x_0}(x_n) \rightarrow x'_0$,
- (P₃) for each strictly positive, continuous linear functional h of X , h_{x_0} exposes x'_0 in B and $h_{x_0}(x'_0) = 0$.

The statement (P₁) is true because $x'_0 \leq P_{x_0}(x) \leq x$ implies $f(x - x'_0) = 0$ hence $x = x_0$.

Definition 3.1. A normed space X ordered by the cone P has the continuous projection property (C.P.P.), if $x \in \text{EP}(P)$ implies that x has a continuous projection.

Proposition 3.2. Let X be a normed space ordered by the cone P . Then:

- (i) X has the C.P.P. if, and only if, $Y = P - P$ has the C.P.P.,
- (ii) if X is a locally solid linear lattice then X has the C.P.P.,
- (iii) if X is a Banach space, X has the R.D.P. and the cone P is closed and generating, then X has the C.P.P.

Proof. If X has the C.P.P. then Y , ordered by the cone P , has the C.P.P. Let Y have the C.P.P. If $x_0 \in \text{EP}(P)$, there exists a continuous positive projection $P_{x_0}(x) = \rho(x_0, x)x_0$ defined on Y . Let $\rho'(x_0, \cdot)$ be a Hahn-Banach extension of $\rho(x_0, \cdot)$ on X . Then $\rho'(x_0, \cdot)$, is positive and $P'_{x_0}(x) = \rho'(x_0, x)x_0$ is a continuous projection of x_0 defined on X . Hence the statement (i) is true. To prove (ii) and (iii) we assume that the cone P is closed and generating and that X has the R.D.P. Since P is closed X is Archimedean. If $x_0 \in \text{EP}(P)$, by [8, Theorem 1.2], there exists a positive linear functional f of X such that

$$f(x) = \sup \{t \in \mathbb{R}_+ \mid tx_0 \leq x\}, \quad \forall x \in P.$$

(In [8, Theorem 1.2], the existence of f is deduced from the fact that X is Archimedean and X has the R.D.P.).

Let $P_{x_0}(x) = f(x)x_0$, $\forall x \in X$. Then P_{x_0} is a linear positive projection and

$P_{x_0}(x) \leq x \forall x \in P$ because the cone P is closed. To show that P_{x_0} is continuous it is enough to show that f is continuous. By [3, 3.5.6.], the statement (iii) is true.

If X is a locally solid linear lattice then X has the R.D.P. and the cone P is generating and closed. Since $0 \leq f(x)x_0 \leq x, \forall x \in P$, there exists $a \in \mathbb{R}_+$ such that

$$|f(x)| \leq \frac{a}{\|x_0\|} \|x\|, \quad \forall x \in P.$$

If $x \in X$ then

$$|f(x)| \leq |f(x^+)| + |f(x^-)| \leq \frac{a}{\|x_0\|} (\|x^+\| + \|x^-\|) \leq \frac{2a^2}{\|x_0\|} \|x\| \leq \frac{2a^3}{\|x_0\|} \|x\|.$$

Hence the statement (ii) is true.

Lemma 3.1. *Let X be a normed space ordered by the cone P , B a base for P defined by $f \in X^*$, $Y = \mathbb{R} \times X$ be ordered by the cone $Y_+ = \mathbb{R}_+ \times P$ and B' be the base for Y_+ defined by the functional $f'(\zeta, x) = \zeta + f(x)$. Then*

- (i) *each extremal point $(\zeta, 0)$ of Y_+ has a continuous projection,*
- (ii) *$x_0 \in \text{sep}(B)$ if, and only if, $(0, x_0) \in \text{sep}(B')$,*
- (iii) *for each $a \in \mathbb{R}_+ \setminus \{0\}$ and $g \in X^*$ we have: g is a negative-strongly exposing functional of x_0 in B if, and only if, $g'(\zeta, x) = -a\zeta + g(x)$ is a negative-strongly exposing functional of $(0, x_0)$ in B' .*

Proof. It is clear that the statement (i) is true.

Let $x_0 \in \text{sep}(B)$ and $g \in X^*$ be a negative-strongly exposing functional of x_0 in B . Then for each $a \in \mathbb{R}_+ \setminus \{0\}$, the functional $g'(\zeta, x) = -a\zeta + g(x)$, exposes $(0, x_0)$ in B' and $g'(0, x_0) = 0$. Let $(\zeta_v, x_v) \in B'$ be such that $g'(\zeta_v, x_v) = -a\zeta_v + g(x_v) \rightarrow 0$. Since $g(x_v) \leq 0$ we have that

$$\zeta_v \rightarrow 0 \quad \text{and} \quad g(x_v) \rightarrow 0.$$

Since $f'(\zeta_v, x_v) = \zeta_v + f(x_v) = 1$ we have that

$$f(x_v) \rightarrow 1.$$

Then $g(x_v/(f(x_v))) \rightarrow 0$, hence $x_v \rightarrow x_0$ and $(\zeta_v, x_v) \rightarrow (0, x_0)$. So $(0, x_0) \in \text{sep}(B')$ and g' is a negative-strongly exposing functional of $(0, x_0)$ in B' .

Let $(0, x_0) \in \text{sep}(B')$ and $h \in Y^*$ be a negative-strongly exposing functional of $(0, x_0)$ in B' . Then there exist $a \in \mathbb{R}_+ \setminus \{0\}$ and $g \in X^*$ such that $h(\zeta, x) = -a\zeta + g(x)$. It is clear that $g(x) \leq 0 \forall x \in P$. If $x_v \in B$ and $g(x_v) \rightarrow g(x_0) = 0$, then $(0, x_v) \in B'$ and $h(0, x_v) = g(x_v) \rightarrow 0 = h(0, x_0)$, hence $x_v \rightarrow x_0$. So the statements (ii) and (iii) are true.

Theorem 3.1. *Let X be a normed space ordered by the cone P , B be a base for P*

defined by $f \in X^*$, $x_0 \in \text{sep}(B)$, g a negative-strongly exposing functional of x_0 in B and $h = f - g$. Then:

- (i) if B' is a base for P defined by $f' \in X^*$, and y_0 is an extreme point of B' with continuous projection P_{y_0} , then $y_0 \in \text{sep}(B')$ and h_{y_0} strongly exposes y_0 in B' .
- (ii) the functional h is uniformly monotonic and the cone P is well-based.

Proof. Since g is a negative-strongly exposing functional of x_0 , we have that $-g$ is positive. Hence h is strictly positive.

Proof of (i). By (P_3) , h_{y_0} exposes y_0 in B' and $h_{y_0}(y_0) = 0$. Let (x_v) be a sequence of B' such that

$$h_{y_0}(x_v) = h(P_{y_0}(x_v) - x_v) \rightarrow 0.$$

If $y_v = x_v - P_{y_0}(x_v)$, then $y_v \in P$ and

$$-h_{y_0}(x_v) = h(y_v) = f(y_v) - g(y_v) \rightarrow 0.$$

Since $f(y_v), -g(y_v) \geq 0$, we have that

$$f(y_v) \rightarrow 0 \quad \text{and} \quad g(y_v) \rightarrow 0.$$

We put

$$z_v = (1 - f(y_v))x_0 + y_v.$$

Then there exists $v_0 \in \mathbb{N}$ such that $z_v \in B$ for each $v \geq v_0$. Since $g(z_v) = g(y_v) \rightarrow 0$ we have that $z_v \rightarrow x_0$, hence $y_v \rightarrow 0$. Since $f'(x_v) = f'(y_v) + f'(P_{y_0}(x_v)) = 1$, we have that $f'(P_{y_0}(x_v)) \rightarrow 1$, hence, by (P_2) , $P_{y_0}(x_v) \rightarrow y_0$. So $x_v = P_{y_0}(x_v) + y_v \rightarrow y_0$, hence h_{y_0} strongly exposes y_0 in B' .

Proof of (ii). Let $Y = \mathbb{R} \times X$ be ordered by the cone $Y_+ = \mathbb{R}_+ \times P$ and B'' be the base for Y_+ defined by the functional $f''(\xi, x) = \xi + f(x)$. Then $(0, x_0) \in \text{sep}(B'')$ and the functional $g'(\xi, x) = -\xi + g(x)$ is a negative-strongly exposing functional of $(0, x_0)$ in B'' .

If $h' = f'' - g'$, then h' is strictly positive and

$$h'(\xi, x) = 2\xi + h(x).$$

Let

$$C = \{(\xi, x) \in Y_+ \mid h'(\xi, x) = 1\}.$$

Then $z_0 = (1/2, 0)$ is an extreme point of C with continuous projection $P_{z_0}(\xi, x) = (\xi, 0)$.

By (i), h'_{z_0} strongly exposes z_0 in C . Moreover, for each $(\xi, x) \in C$ we have

$$|h'_{z_0}(\xi, x)| \leq h'(\xi, x) = 1.$$

By Proposition 3.1, C is bounded. So the base for P defined by h is bounded. Hence P is well-based and the functional h is uniformly monotonic.

Proposition 3.3. *Let X be a normed space ordered by the cone P and B be a base for P defined by $f \in X^*$. If h is a continuous uniformly monotonic linear functional of X , then for each extreme point x_0 of B with continuous projection P_{x_0} , the functional h_{x_0} strongly exposes x_0 in B .*

Proof. Since h is uniformly monotonic, there exists $a \in \mathbb{R}_+ \setminus \{0\}$ such that $h(x) \geq a\|x\|$, for each $x \in P$. Let $C = \{x \in P \mid h(x) = 1\}$ and $x'_0 = \lambda x_0 \in C$. To prove that h_{x_0} strongly exposes x_0 in B , by Theorem 3.1, it is enough to prove that h_{x_0} strongly exposes x'_0 in C . Now h_{x_0} exposes x'_0 in C and $h_{x_0}(x'_0) = 0$. Let (x_v) be a sequence of C such that

$$h_{x_0}(x_v) = h(P_{x_0}(x_v) - x_v) \rightarrow 0.$$

Then $h(P_{x_0}(x_v)) \rightarrow 1$, hence

$$P_{x_0}(x_v) \rightarrow x'_0.$$

If $y_v = x_v - P_{x_0}(x_v)$, then $y_v \in P$ and

$$h(x_v) = h(P_{x_0}(x_v)) + h(y_v) = 1,$$

hence $h(y_v) \rightarrow 0$. So we have that $y_v \rightarrow 0$, because $a\|y_v\| \leq h(y_v)$. Hence $x_v \rightarrow x'_0$.

Corollary 3.1. *Let X be a Banach space ordered by the closed, generating cone P , B be a base for P and X have the R.D.P. If $\text{ep}(B) \neq \emptyset$ then the following statements are equivalent:*

- (i) $\text{sep}(B) \neq \emptyset$,
- (ii) $\text{ep}(B) = \text{sep}(B)$,
- (iii) P is well-based.

Proof. By Proposition 3.2 X has the C.P.P. Also by [3, 3.5.6], each base for P is defined by a continuous linear functional. By Theorem 3.1 we have that (i) \Rightarrow (ii) and (i) \Rightarrow (iii). It is clear that (ii) \Rightarrow (i). If the cone P is well-based, by [3, 3.8.12], there exists a uniformly monotonic, continuous linear functional of X and by Proposition 3.3 we have that (iii) \Rightarrow (i).

Proposition 3.4. *Let X be a normed space ordered by the cone P and B be a base for P defined by $f \in X^*$. If x_0 is an extreme point of B with continuous projection P_{x_0} , then*

- (i) g is a negative-strongly exposing functional of x_0 in B if, and only if, there exists a uniformly monotonic, continuous linear functional h of X such that $g = h(x_0)\rho(x_0, \cdot) - h$.
- (ii) g strongly exposes x_0 in B if, and only if, there exists a uniformly monotonic, continuous, linear functional h of X and $\lambda \in \mathbb{R}$ such that $g = h(x_0)\rho(x_0, \cdot) - h + \lambda f$.

Proof. Let $h \in X^*$ be uniformly monotonic. By Proposition 3.3, $g = h_{x_0} = h(x_0)\rho(x_0, \cdot) - h$ is a negative-strongly exposing functional of x_0 in B . Let g strongly exposes x_0 in B and $g(x_0) = 0$. By Theorem 3.1, $h = f - g$ is uniformly monotonic. Let

$$w = h - f + \rho(x_0, \cdot).$$

Then $w(x_0) = 1$ because $f(x_0) - h(x_0) = g(x_0) = 0$. So

$$g = w(x_0)\rho(x_0, \cdot) - w.$$

To show that w is uniformly monotonic it is enough to show that $w(x) \geq \gamma > 0$, $\forall x \in B' = \{x \in P \mid h(x) = 1\}$, because then $w(x) \geq \gamma h(x) \geq \gamma a \|x\| \forall x \in P$. Let $w(x_v) \rightarrow 0$ for a sequence (x_v) of B' . Then $w(x_v) = -g(x_v) + \rho(x_0, x_v) \rightarrow 0$ and therefore

$$g(x_v) \rightarrow 0 \quad \text{and} \quad \rho(x_0, x_v) \rightarrow 0.$$

Moreover, $f(x_v) \rightarrow 1$ because $h(x_v) = f(x_v) - g(x_v) = 1$. If $y_v = x_v / (f(x_v))$ then $y_v \in B$ and $g(y_v) \rightarrow 0$. So $y_v \rightarrow x_0$, hence $x_v \rightarrow x_0$ and therefore $\rho(x_0, x_v) \rightarrow 1$. This is a contradiction, hence w is uniformly monotonic and the proof of (i) is complete. The statement (ii) follows by (i) because g strongly exposes x_0 in B if, and only if, $g + \lambda f$ strongly exposes x_0 in B .

Proposition 3.5. Let X be a normed space ordered by the cone P and B be a base for P defined by $f \in X^*$. If x_0 is an extreme point of B with continuous projection P_{x_0} then the following statements are equivalent:

- (i) B is bounded,
- (ii) for each sequence (x_v) of B , $x_v \rightarrow x_0$ if, and only if, $P_{x_0}(x_v) \rightarrow x_0$.

Proof. Let B be bounded. Then f is uniformly monotonic and by Proposition 3.3, f_{x_0} strongly exposes x_0 in B . Let (x_v) be a sequence of B . If $x_v \rightarrow x_0$, then $P_{x_0}(x_v) \rightarrow P_{x_0}(x_0) = x_0$. If $P_{x_0}(x_v) \rightarrow x_0$, then

$$f_{x_0}(x_v) = f(P_{x_0}(x_v) - x_v) = f(P_{x_0}(x_v)) - 1 \rightarrow f(x_0) - 1 = 0 = f_{x_0}(x_0).$$

Hence $x_v \rightarrow x_0$. So (i) \Rightarrow (ii).

Let the statement (ii) be true. To show that B is bounded, it is enough to show that the functional f_{x_0} which is bounded on B strongly exposes x_0 in B . (Proposition 3.1.)

Let (x_v) be a sequence of B such that

$$f_{x_0}(x_v) \rightarrow f_{x_0}(x_0) = 0.$$

Then $f_{x_0}(x_v) = f(P_{x_0}(x_v)) - 1 \rightarrow 0$, hence $P_{x_0}(x_v) \rightarrow x_0$ and $x_v \rightarrow x_0$. So f_{x_0} strongly exposes x_0 in B .

4. Characterizations of $l_1(\Gamma)$

Let G be a closed and convex subset of a Banach space X . The set G has the Krein–Milman property (K.M.P.) if $K = \overline{\text{coep}}(K)$, for each closed, convex and bounded subset K of G . It is known, [2, 3.5.7], that the set G has the Radon–Nikodým property (R.N.P.) if, and only if, $K = \overline{\text{cosep}}(K)$, for each closed, convex and bounded subset K of G . Moreover we know, [1], that in locally solid lattice Banach spaces the R.N.P. and the K.M.P. are equivalent. Let Γ be any set. We denote by $l_1(\Gamma)$ the Banach space of all functions $\xi: \Gamma \rightarrow \mathbb{R}$, $\xi = (\xi(i))_{i \in \Gamma}$, such that $\sum_{i \in \Gamma} |\xi(i)| < +\infty$, with norm $\|\xi\| = \sum_{i \in \Gamma} |\xi(i)|$. The space $l_1(\Gamma)$ has the R.N.P., [2, 4.1.9] and ordered by the cone $l_1^+(\Gamma) = \{\xi \in l_1(\Gamma) \mid \xi(i) \geq 0 \forall i \in \Gamma\}$ is a Banach lattice. The set $B = \{\xi \in l_1^+(\Gamma) \mid \|\xi\| = 1\}$ is a closed bounded base for the cone $l_1^+(\Gamma)$. We denote by l_1 space $l_1(\mathbb{N})$.

Theorem 4.1. *Let X be an infinite-dimensional Banach space ordered by the closed, generating cone P and X have the R.D.P. Then:*

- (i) *X is order-isomorphic to $l_1(\Gamma)$ if, and only if, P has a closed, bounded base with the K.M.P.;*
- (ii) *X is order-isomorphic to l_1 if, and only if, P has a separable, closed, bounded base with the K.M.P.*

Proof. Let T be an order-isomorphism of X onto $l_1(\Gamma)$. Since $B = \{\xi \in l_1^+(\Gamma) \mid \|\xi\| = 1\}$ is a closed and bounded base for $l_1^+(\Gamma)$ with the K.M.P. we have that $T^{-1}(B)$ is a closed, bounded base for P with the K.M.P. Let B be a closed, bounded base for P defined by the functional f and let B have the K.M.P. By [3, 3.5.6], the functional f is continuous. Let

$$\text{ep}(B) = \{b_i \mid i \in \Gamma\}.$$

By Proposition 3.2 we have that X has the C.P.P.

We shall prove that

$$x = \sum_{i \in \Gamma} \rho(b_i, x) b_i \quad \text{and} \quad \sum_{i \in \Gamma} \rho(b_i, x) < +\infty, \quad \forall x \in P. \quad (1)$$

At first we shall show that

$$L = \{x \in P \mid P_{b_i}(x) = 0, \quad \forall i \in \Gamma\} = \{0\}.$$

The set L is a cone and it is closed because P_{b_i} is continuous $\forall i \in \Gamma$. If $L \neq \{0\}$, the set $B' = B \cap L$ is a non-empty, closed and bounded base for L . So $\text{EP}(L) \neq \emptyset$ because $\text{ep}(B') \neq \emptyset$. Also $\text{EP}(L) \subseteq \text{EP}(P)$ because for each $x \in L$ and $y \in P$, $0 \leq y \leq x$ implies that $y \in L$. Hence $b_j \in \text{EP}(L)$ for at least one $j \in \Gamma$. This contradicts the definition of L because $P_{b_j}(b_j) = b_j$. Hence $L = \{0\}$.

We denote by F the set of finite subsets of Γ and for each $x \in P$ and $\delta \in F$ we denote by x_δ the sum

$$x_\delta = \sum_{i \in \delta} P_{b_i}(x).$$

Let $x \in B$. Then $(x_\delta)_{\delta \in F}$ is an upward-directed net of P (if $\delta_1, \delta_2 \in F$ we say that $\delta_1 \leq \delta_2$ if, and only if, $\delta_1 \subseteq \delta_2$). We shall show that $x_\delta = \sup\{P_{b_i}(x) \mid i \in \delta\}$. If $z \geq P_{b_i}(x) \forall i \in \delta$, then $w = z - P_{b_i}(x) \geq 0$. If $j \in \delta$ and $j \neq i$, then $w \geq P_{b_j}(w) = P_{b_j}(z) \geq P_{b_j}(x)$ and therefore $z \geq P_{b_i}(x) + P_{b_j}(x)$. By a similar process we have that $z \geq x_\delta$, hence

$$x_\delta = \sup\{P_{b_i}(x) \mid i \in \delta\} \leq x, \quad \forall \delta \in F.$$

By [3, 3.8.8], we have

$$y = \lim x_\delta = \sup_{\delta \in F} (x_\delta) \leq x.$$

This implies that $P_{b_i}(x) \leq y \leq x \forall i \in \Gamma$ and therefore that $P_{b_i}(x - y) = 0 \forall i \in \Gamma$ because $P_{b_i}(x) \leq P_{b_i}(y) \leq P_{b_i}(x) \forall i \in \Gamma$. Hence $x = y$ and therefore

$$x = \sum_{i \in \Gamma} \rho(b_i, x) b_i.$$

Since $f \in X^*$ and f define the base B we have that

$$f(x) = \sum_{i \in \Gamma} \rho(b_i, x) = 1.$$

So (1) is true because it is true for each $x \in B$.

We define the map $T: P \rightarrow l_1^+(\Gamma)$ as follows:

$$T(x) = (\rho(b_i, x))_{i \in \Gamma}, \quad \forall x \in P.$$

It is clear that $T(\lambda x + \mu y) = \lambda T(x) + \mu T(y)$, $\forall x, y \in P$ and $\lambda, \mu \in \mathbb{R}_+$. T is one-to-one because

$$x = \sum_{i \in \Gamma} \rho(b_i, x) b_i, \quad \forall x \in P.$$

Since the set B is bounded there exists $M \in \mathbb{R}_+$ such that

$$\|x\| \leq M, \quad \forall x \in B.$$

We shall show that the map T is onto.

Let $\xi = (\xi(i))_{i \in \Gamma} \in l_1^+(\Gamma)$. For each $\delta \in F$ we put

$$x_\delta = \sum_{i \in \delta} \xi(i) b_i \quad \text{and} \quad \xi_\delta = T(x_\delta).$$

Let $\varepsilon > 0$. Since $\lim \xi_\delta = \xi$ there exists $\delta_0 \in F$ such that

$$\|\xi_{\delta_1} - \xi_{\delta_2}\| < \varepsilon, \quad \forall \delta_1, \delta_2 > \delta_0.$$

If $\delta'_1 = \delta_1 \setminus \delta_2$ and $\delta'_2 = \delta_2 \setminus \delta_1$ then $\xi_{\delta'_1} - \xi_{\delta'_2} = \xi_{\delta_1} - \xi_{\delta_2}$. Since $\xi_{\delta'_1}, \xi_{\delta'_2}$ are disjoint we have that

$$|\xi_{\delta'_1} - \xi_{\delta'_2}| = |\xi_{\delta_1} - \xi_{\delta_2}| = \xi_{\delta_1} + \xi_{\delta_2}$$

and therefore

$$\|\xi_{\delta_1} - \xi_{\delta_2}\| = \|\xi_{\delta'_1} + \xi_{\delta'_2}\| = \|\xi_{\delta'_1}\| + \|\xi_{\delta'_2}\| \leq \varepsilon.$$

So

$$\|x_{\delta_1} - x_{\delta_2}\| = \|x_{\delta'_1} - x_{\delta'_2}\| \leq M(\|\xi_{\delta'_1}\| + \|\xi_{\delta'_2}\|) < M\varepsilon.$$

Hence the net $(x_\delta)_{\delta \in F}$ is Cauchy. If $x = \lim x_\delta$, then $x \in P$ and $\rho(b_i, x) = \xi(i) \forall i \in \Gamma$. So $T(x) = \xi$ and the map T is onto $l_1^+(\Gamma)$. By [3, 1.5.6] T can be extended to a linear map T of X into $l_1(\Gamma)$ as follows: $T(x) = T(y) - T(z)$, where $x = y - z$ and $y, z \in P$. Since $l_1^+(\Gamma)$ is generating the map T is onto $l_1(\Gamma)$. Also T is one-to-one because $T(x) = 0$ implies $T(y) = T(z)$ and therefore $y = z$. By the definition of T we have that T and T^{-1} are positive.

Let $\xi = T(x) = (\rho(b_i, x))_{i \in \Gamma} \in l_1^+(\Gamma)$. Then

$$\|T^{-1}(\xi)\| = \|x\| = \left\| \sum_{i \in \Gamma} \rho(b_i, x) b_i \right\| \leq M \sum_{i \in \Gamma} \rho(b_i, x) = M \|\xi\|.$$

Since the map T^{-1} is linear and $l_1(\Gamma)$ is a Banach lattice we have that T^{-1} is continuous. By the open mapping theorem, T is continuous. So T is an order-isomorphism of X onto $l_1(\Gamma)$ and the statement (i) is true.

Let B be a separable, closed and bounded base for P and let B have the K.M.P. Then X is order-isomorphic to $l_1(\Gamma)$ and the base $C = T(B)$ for $l_1^+(\Gamma)$ is separable. Also there exists $\lambda \in \mathbb{R}_+$ such that $\|\xi\| \geq \lambda > 0$ for each $\xi \in C$. Let $\text{ep}(C) = \{\xi_i \mid i \in \Gamma\}$. Then

$$\|\xi_i - \xi_j\| = \|\xi_i\| + \|\xi_j\| \geq 2\lambda, \quad \forall i \neq j.$$

So the set Γ is countable because C is separable and $\|\xi_i - \xi_j\| \geq 2\lambda, \forall i \neq j$. Hence X is ordered isomorphic to l_1 and the statement (ii) is true.

Let K be a closed, convex, unbounded subset of a Banach space X . For each real number $\rho > 0$ we denote by $K_\rho, K_{s, \rho}$, the sets

$$\{x \in K \mid \|x\| \leq \rho\}, \{x \in K \mid \|x\| = \rho\}$$

respectively, whenever these sets are non-empty. In [7, Propositions 1 and 3] it is proved:

- (i) if X has the K.M.P. then $K_\rho \neq \overline{\text{co}} K_{s, \rho}$ for at least one $\rho \in \mathbb{R}_+$ implies $\text{ep}(K) \neq \emptyset$;

(ii) if X has the R.N.P., then: $K_\rho \neq \overline{\text{co}}K_{s,\rho}$ for at least one $\rho \in \mathbb{R}_+ \Leftrightarrow K$ is dentable $\Leftrightarrow \text{sep}(K) \neq \emptyset$.

It is easy to show that the proof of these results can be accepted for the case where K is a subset of a closed, convex and unbounded subset A of X and A has the K.M.P., the R.N.P. respectively.

In [7, Corollary 3], it is shown that each closed and convex subset of l_1^+ has at least one strongly exposed point. In the following proposition we prove a similar result for well-based cones.

Proposition 4.1. *Let X be a Banach space ordered by the closed, well-based cone P and K a closed and convex subset of P . If P has the K.M.P. (respectively, the R.N.P.) then $\text{ep}(K) \neq \emptyset$ (respectively, $\text{sep}(K) \neq \emptyset$).*

Proof. If the set K is bounded the proposition is true. Let K be unbounded. To show that $\text{ep}(K) \neq \emptyset$ (respectively, $\text{sep}(K) \neq \emptyset$) it is enough to show that $\overline{\text{co}}K_{s,\rho} \neq K_\rho$, for at least one $\rho \in \mathbb{R}_+$. Let a uniformly monotonic, continuous linear functional f of X ($f(x) \geq a\|x\|$) $\forall x \in P$, $x_0 \in K$ and a real number $\varepsilon > 0$. If $\rho > \|x_0\|$ and $a\rho > f(x_0) + \varepsilon$, then $x_0 \in K_\rho$ and for each convex combination $x = \sum_{i=1}^n \lambda_i x_i$ of elements of $K_{s,\rho}$ we have

$$f(x) = \sum_{i=1}^n \lambda_i f(x_i) \geq \sum_{i=1}^n \lambda_i a\|x_i\| = a\rho > f(x_0) + \varepsilon.$$

Hence for each $y \in \overline{\text{co}}K_{s,\rho}$ we have that

$$f(y) \geq f(x_0) + \varepsilon > f(x_0)$$

and therefore $\overline{\text{co}}K_{s,\rho} \neq K_\rho$.

Proposition 4.2. *Let X be an infinite-dimensional Banach space ordered by the closed, generating cone P and X have the R.D.P.*

If P has the K.M.P. the statements (i), (ii), (iii), (iv) and (v) are equivalent.

If P has the R.N.P. all the following statements are equivalent:

- (i) X is order-isomorphic to $l_1(\Gamma)$,
- (ii) P is well-based,
- (iii) $\text{sep}(B) \neq \emptyset$, for at least one base B for P ,
- (iv) $0 \in \text{sep}(P)$,
- (v) $\text{sep}(K) \neq \emptyset$ for each closed and convex subset K of P ,
- (vi) B is dentable, for at least one base B for P ,
- (vii) P is dentable,
- (viii) K is dentable, for each closed and convex subset K of P .

Proof. Let P have the K.M.P. If P is well-based, there exists a uniformly monotonic continuous linear functional f of X . This functional defines a closed and bounded base

C for P and C has the K.M.P. Hence, by Theorem 4.1, (ii) \Leftrightarrow (i). Since $\text{ep}(C) \neq \emptyset$, by Corollary 3.1, we have that (ii) \Leftrightarrow (iii). It is easy to show that (ii) \Leftrightarrow (iv). Also (v) \Rightarrow (iii) \Rightarrow (ii). Since (ii) \Leftrightarrow (i), the statement (ii) implies that P has the R.N.P. and by Proposition 4.1 we have that (ii) \Rightarrow (v).

Let P have the R.N.P. Then for each closed and convex subset A of P we have: $\text{sep}(A) \neq \emptyset$ if and only if, A is dentable. Hence (iii) \Leftrightarrow (vi), (iv) \Leftrightarrow (vii) and (v) \Leftrightarrow (viii).

REFERENCES

1. J. BOURGAIN and M. TALAGRAND, Dans un espace de Banach reticulé solid, la propriété de Radon–Nikodým et celle de Krein–Milman sont équivalentes, *Proc. Amer. Math. Soc.* **81** (1981), 93–96.
2. R. D. BOURGIN, *Geometric Aspects of Convex Sets with the Radon–Nikodým Property* (Lecture Notes in Mathematics 993).
3. G. J. O. JAMESON, *Ordered Linear Spaces* (Lecture Notes in Mathematics, 141).
4. D. KENDALL, Simplexes and vector lattices, *J. London Math. Soc.* **37** (1962), 365–371.
5. R. R. PHELPS, Dentability and extreme points in Banach spaces, *J. Functional Analysis* **16** (1974), 78–90.
6. I. A. POLYRAKIS, Lattice Banach spaces order-isomorphic to l_1 , *Math. Proc. Cambridge Phil. Soc.* **94** (1983), 519–522.
7. I. A. POLYRAKIS, Extreme points of unbounded, closed and convex sets in Banach spaces, *Math. Proc. Cambridge Phil. Soc.* **95** (1984), 319–323.
8. G. C. SCHMIDT, Extensions theorems for linear lattices with positive algebraic basis, *Periodica Mathematica Hungarica* **6**(4), (1975), 295–307.

DEPARTMENT OF MATHEMATICS
NATIONAL TECHNICAL UNIVERSITY
ZOGRAFON CAMPUS, 157 73
ATHENS, GREECE